



# The Journal of Biomedical Research

## Genetic variants in the Hedgehog signaling pathway genes are associated with gastric cancer risk in a Chinese Han population

Zhang Yujuan, Lu Kai, Wu Xu, Liu Hanting, Xin Junyi, Wang Xiaowei, Gong Weida, Zhao Qinghong, Wang Meilin, Chu Haiyan, Du Mulong, Tao Guoquan, Zhang Zhengdong

Cite this article as:

Zhang Yujuan, Lu Kai, Wu Xu, Liu Hanting, Xin Junyi, Wang Xiaowei, Gong Weida, Zhao Qinghong, Wang Meilin, Chu Haiyan, Du Mulong, Tao Guoquan, Zhang Zhengdong. Genetic variants in the Hedgehog signaling pathway genes are associated with gastric cancer risk in a Chinese Han population[J]. *Journal of Biomedical Research*, 2022, 36(1): 22–31. doi: 10.7555/JBR.35.20210091

View online: <https://doi.org/10.7555/JBR.35.20210091>

---

### Articles you may be interested in

[Dietary vitamin D intake and vitamin D related genetic polymorphisms are not associated with gastric cancer in a hospital-based case-control study in Korea](#)

The Journal of Biomedical Research. 2018, 32(4): 257 <https://doi.org/10.7555/JBR.32.20170089>

[Activation of p38/HSP27 pathway counters melatonin-induced inhibitory effect on proliferation of human gastric cancer cells](#)

The Journal of Biomedical Research. 2019, 33(5): 317 <https://doi.org/10.7555/JBR.33.20180066>

[A novel long non-coding RNA NFIA-AS1 is down-regulated in gastric cancer and inhibits proliferation of gastric cancer cells](#)

The Journal of Biomedical Research. 2019, 33(6): 371 <https://doi.org/10.7555/JBR.33.20190015>

[Superior \*in vitro\* anticancer effect of biomimetic paclitaxel and triptolide co-delivery system in gastric cancer](#)

The Journal of Biomedical Research. 2021, 35(4): 327 <https://doi.org/10.7555/JBR.35.20210102>

[miR-3622b-5p regulates cisplatin resistance of human gastric cancer cell line by targeting BIRC5](#)

The Journal of Biomedical Research. 2019, 33(6): 382 <https://doi.org/10.7555/JBR.33.20180078>

[Association of long non-coding RNA HOTAIR and MALAT1 variants with cervical cancer risk in Han Chinese women](#)

The Journal of Biomedical Research. 2019, 33(5): 308 <https://doi.org/10.7555/JBR.33.20180096>



# Genetic variants in the Hedgehog signaling pathway genes are associated with gastric cancer risk in a Chinese Han population

Yujuan Zhang<sup>1,2,Δ</sup>, Kai Lu<sup>1,2,Δ</sup>, Xu Wu<sup>3,Δ</sup>, Hanting Liu<sup>1,2</sup>, Junyi Xin<sup>1,2</sup>, Xiaowei Wang<sup>1,2</sup>, Weida Gong<sup>4</sup>, Qinghong Zhao<sup>5</sup>, Meilin Wang<sup>1,2,6</sup>, Haiyan Chu<sup>1,2</sup>, Mulong Du<sup>1,7,✉</sup>, Guoquan Tao<sup>3,✉</sup>, Zhengdong Zhang<sup>1,2,6,✉</sup>

<sup>1</sup>Department of Environmental Genomics, Jiangsu Key Laboratory of Cancer Biomarkers, Prevention and Treatment, Collaborative Innovation Center for Cancer Personalized Medicine, Nanjing Medical University, Nanjing, Jiangsu 211166, China;

<sup>2</sup>Department of Genetic Toxicology, the Key Laboratory of Modern Toxicology of Ministry of Education, Center for Global Health, School of Public Health, Nanjing Medical University, Nanjing, Jiangsu 211166, China;

<sup>3</sup>Department of General Surgery, the Affiliated Huai'an No.1 People's Hospital of Nanjing Medical University, Huai'an, Jiangsu 223300, China;

<sup>4</sup>Department of General Surgery, Yixing People's Hospital, Yixing, Jiangsu 214200, China;

<sup>5</sup>Department of General Surgery, the Second Affiliated Hospital of Nanjing Medical University, Nanjing, Jiangsu 210011, China;

<sup>6</sup>Department of Environmental Genomics, Center for Global Health, School of Public Health, Nanjing Medical University, Nanjing, Jiangsu 211166, China;

<sup>7</sup>Department of Biostatistics, Center for Global Health, School of Public Health, Nanjing Medical University, Nanjing 211166, China.

## Abstract

The Hedgehog signaling pathway participates in the occurrence and progression of cancers including gastric cancer. We conducted this study to evaluate whether genetic variants in the Hedgehog signaling pathway genes would affect gastric cancer risk. Multi-marker Analysis of GenoMic Annotation (MAGMA) was used to investigate the aggregated genetic effects of single nucleotide polymorphisms (SNPs) assigned to candidate genes. The relationship between SNPs and gastric cancer risk was estimated by multivariate logistic regression analyses. Gene expression was calculated using databases obtained from The Cancer Genome Atlas (TCGA) and The Gene Expression Omnibus (GEO). Kaplan - Meier plotter was used to evaluate the association between gene expression with gastric cancer survival. Tumor Immune Estimation Resource 2.0 (TIMER 2.0) was applied to determine the correlation between selected gene expression and the immune cell infiltration degree. We identified that the G allele of rs2990912 in *KIF27* was associated with higher gastric cancer risk, especially in the young and male subgroups. The expression of *KIF27* in gastric cancer tissues was higher than that in normal tissues, leading to poor survival in gastric cancer patients. Besides, *KIF27* expression was related to immune cell infiltration and positively correlated with PD-L1 expression. Our findings highlight the key role of genetic variation in the

<sup>Δ</sup>These authors contributed equally to this work.

<sup>✉</sup>Corresponding authors: Mulong Du, Department of Biostatistics, School of Public Health, Nanjing Medical University, 101 Longmian Avenue, Jiangning District, Nanjing, Jiangsu 211166, China. Tel: +86-25-86868423, E-mail: [drdumulong@njmu.edu.cn](mailto:drdumulong@njmu.edu.cn); Guoquan Tao, Department of General Surgery, the Affiliated Huai'an No.1 People's Hospital of Nanjing Medical University, No. 1 Huanghe Western Road, Huaiyin District, Huai'an, Jiangsu 223300, China. Tel: +86-517-84922412, E-mail: [taoguoquan5698102@163.com](mailto:taoguoquan5698102@163.com); Zhengdong Zhang, School of Public Health, Nanjing Medical University, 101 Longmian Avenue,

Jiangning District, Nanjing, Jiangsu 211166, China. Tel/Fax: +86-25-86868423/+86-25-86868499, E-mail: [drzdzhang@njmu.edu.cn](mailto:drzdzhang@njmu.edu.cn).

Received: 09 June 2021; Revised: 30 August 2021; Accepted: 06 September 2021; Published online: 29 October 2021

CLC number: R735.2, Document code: A

The authors reported no conflict of interests.

This is an open access article under the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited.

Hedgehog signaling pathway genes in gastric cancer susceptibility, which may provide important insights into the diagnosis, prognosis, and treatment of gastric cancer.

**Keywords:** gastric cancer, Hedgehog signaling pathway, genetic susceptibility, molecular epidemiology

---

## Introduction

Gastric cancer is the fifth most common cancer globally, with more than 1 000 000 new cases are diagnosed each year and is responsible for more than 8% of all oncological deaths in 2018<sup>[1]</sup>. The incidence of gastric cancer is the highest in East Asia, especially in China, followed by Central and Eastern Europe<sup>[1]</sup>. Gastric cancer ranks the second in incidence and mortality in China<sup>[2]</sup>. Although there is a steady decreasing trend in the incidence and mortality rates of gastric cancer because of the change of dietary habits, decrease in *Helicobacter pylori* infection, and reduction in the smoking prevalence, gastric cancer is still a huge threat to human health due to its nonspecific early stage disease symptomatology and the lack of effective screening methods<sup>[3]</sup>. More than 80% of patients with gastric cancer are diagnosed as advanced with poor clinical outcomes and the 5-year survival rate is less than 20%<sup>[4]</sup>. Therefore, studies on the mechanism of gastric cancer occurrence and development may improve the early diagnosis and prognosis of gastric cancer.

Genome-wide association studies (GWASs) have evaluated a large number of single-nucleotide polymorphisms (SNPs) related to complex diseases, including gastric cancer<sup>[5–6]</sup>. The standard approach of GWAS in single-variant level misses an amount of gastric cancer risk-related SNPs with a moderate effect because of the strict threshold at  $P < 5 \times 10^{-8}$ . Gene-based and pathway-based analyses have been applied in GWAS to identify SNPs with moderate effects but significantly related to disease by adopting a looser threshold<sup>[7–9]</sup>.

The Hedgehog signaling pathway is an evolutionary conservative pathway that transmits signals from the cell membrane. Its members have been considered to participate in embryonic development, adult tissue homeostasis, tissue repair, and carcinogenesis<sup>[10–11]</sup>. The Hedgehog signaling pathway mainly consists of the Hedgehog (HH) ligands, which include Sonic hedgehog (SHH), Indian hedgehog (IHH) and Desert hedgehog (DHH), Smoothed (SMO), Patched 1 (PTCH1), Patched 2 (PTCH2), Suppressor of Fused (SUFU), Kinesin family member 7 (KIF7), Kinesin

family member 27 (KIF27), GLI family zinc finger 1 (GLI1), GLI family zinc finger 2 (GLI2), GLI family zinc finger 3 (GLI3), and GLI family zinc finger 4 (GLI4)<sup>[12]</sup>. This pathway exerts its biological function by affecting the balance between GLI activator form (GLIA) and GLI repressor form (GLIR). The Hedgehog signaling pathway is associated with self-renewal, survival, proliferation, and invasion of tumor cells and related to angiogenesis, fibrosis, immune evasion, and neuropathic pain in the tumor microenvironment<sup>[13]</sup>. GLI2 overexpression induces loss of gastrin and antral hyperplasia<sup>[14]</sup>. A lower Hedgehog-interacting protein (HHIP) level is positively correlated with gastric cancer metastasis. HHIP can mediate the progression and metastasis of gastric cancer by regulating its promoter methylation level in a feedback manner<sup>[15]</sup>. miRNA-150 targets *Sufu*, a tumor suppressor, and subsequently promotes cell proliferation, migration, and epithelial-mesenchymal transition *via* dual activation of Hedgehog and Wnt/ $\beta$ -catenin signaling pathway<sup>[16]</sup>. GLI mediates mTOR-induced programmed death-ligand 1 (PD-L1) expression in gastric cancer<sup>[17]</sup>. Apatinib, an oral small molecule tyrosine kinase inhibitor used to treat advanced gastric cancer, inhibits gastric cancer stem cells by blocking the SHH pathway<sup>[18]</sup>. Several studies have reported that genetic variants in the Hedgehog signaling pathway genes can influence the risk of various cancers such as bladder cancer<sup>[19–20]</sup>, but most of them only concentrate on a few genes, and the existing evidence for the influence of SNPs in the Hedgehog signaling pathway genes on gastric cancer susceptibility in Chinese population is limited.

In this study, we used a comprehensive strategy to screen the effective SNPs in key genes of Hedgehog signaling pathway using bioinformatics tools and functional analysis, and explored the possible mechanisms of the risk SNPs and corresponding host genes involved in gastric oncogenesis.

## Materials and methods

### Study subjects and genotyping

Gastric cancer GWAS data (phs000361. v1. p1)

was downloaded from the database of Genotype and Phenotype. The Illumina 660W Quad chip was used for genotyping. The nongenotyped variants were imputed on the basis of the 1000 Genomes Project using IMPUTE for the gastric cancer GWAS<sup>[21]</sup>. A total of 3725 individuals (1625 gastric cancer cases and 2100 controls) were included in this study. This population was derived from 4987 samples of the case-control and case-only components of the Shanxi Upper Gastrointestinal Cancer Genetics Project (Shanxi). The population characteristics were described previously<sup>[22]</sup> and characteristic information was summarized in **Supplementary Table 1** (available online).


### Steps for selecting genes and SNPs

We selected 31 key Hedgehog signaling pathway genes from Kyoto Encyclopedia of Genes and Genomes (KEGG, <https://www.genome.jp/kegg/>) and the previously published studies. Gene position was obtained from UCSC Genome Browser (**Supplementary Table 2**, available online, <https://genome.ucsc.edu/>). A total of 17 962 SNPs in the 31 reference gene regions were genotyped. **Fig. 1** shows SNPs screening process. Firstly, we determined 3164 eligible SNPs in 27 genes after the quality control using the criteria as: minor allele frequency (MAF) >0.05,  $P_{(HWE)} > 0.05$  for Hardy-Weinberg equilibrium (HWE), and call rate >95%. Then, we evaluated the aggregated single-variant effect into each candidate gene on gastric cancer risk using Multi-marker

Analysis of GenoMic Annotation (MAGMA)<sup>[23]</sup>. MAGMA is a tool for gene-based and gene-set pathway analysis with GWAS data. The Z-value of MAGMA reflects the strength of the association between each gene and the phenotype, and a higher value corresponds to a stronger association. Finally, we predicted the function of SNPs located at risk genes using Regulome DB (<https://regulomedb.org/regulome-search/>), SNP info Web Server (<https://manticore.niehs.nih.gov/>), and HaploReg (<https://pubs.broadinstitute.org/mammals/haploreg/haploreg.php>), and selected the tagged SNPs using pair-wise linkage disequilibrium (LD) analysis based on the GWAS data ( $r^2 \geq 0.8$ ). SNPs with a score higher than 5 in Regulome DB were removed.

### Expression analysis

Gene expression data, genotype data, cancer pathological feature, and *H. pylori* infection information of patients with gastric cancer were downloaded from The Cancer Genome Atlas (TCGA) stomach adenocarcinoma (STAD) datasets. Gene expression information obtained from the Gene Expression Omnibus (GEO) database (GSE66229, GSE13911, and GSE37023) was used to analyze the expression pattern of genes in gastric tumor and normal tissues. The GTEx dataset (<https://www.gtexportal.org/home/>) was used to analyze the expression quantitative trait loci (eQTL) for candidate SNPs on genes. The influence of gene expression on



**Fig. 1 Flow diagram.** Flowchart for selecting SNPs in the Hedgehog signaling pathway. MAF: minor allele frequency; HWE: Hardy-Weinberg equilibrium; LD: linkage disequilibrium; FDR: false discovery rate.

gastric cancer patient's survival was measured by Kaplan-Meier plotter. Correlation analysis was conducted by the Gene Expression Profiling Interactive Analysis (GEPIA, <http://gepia.cancer-pku.cn/>).

### Immune infiltration analysis

Tumor Immune Estimation Resource 2.0 (TIMER2.0) was used to explore the correlation between candidate gene expression and immune cell infiltration degree. This tool is useful for systematic analysis of the immune infiltration across pan cancers, providing the abundance of immune infiltrates estimated by a variety of immune deconvolution methods according to the expression profiles of TCGA tumors. TIMER, xCell, MCP-counter, CIBERSORT, EPIC, and quantTIseq were used to estimate the level of immune infiltration. The "Gene" module was used to analyze the correlation between gene expression and the level of immune infiltration in gastric cancer. The analysis results were visualized by scatter plots. By selecting the "Purity Adjustment" option, partial Spearman correlation was used for correlation analysis, and the Rho value was performed to indicate the degree of correlation. The "Gene\_Corr" module was used to explore the correlation between the selected genes and immune markers in gastric cancer.

### Statistical analysis

The distributions of variables in gastric cancer cases and controls were assessed using a Student's *t*-test. A Chi-square test was performed to analyze HWE in controls. Odds ratios (ORs) and their 95% confidence intervals (CIs) were examined using multivariate logistic regression analyses to evaluate genetic variants of SNPs on gastric cancer risk. The covariates of age and sex were adjusted. False discovery rate (FDR) correction was performed to adjust *P* values. The two-sided Mann-Whitney test was utilized to compare the significance of different gene expression between gastric tumor and normal tissues. R (version 3.5.1) and PLINK (version 1.90) were used to perform all statistical analyses.

## Results

### Gene-set analysis and SNP selection

The mechanism of 31 genes involved in the Hedgehog signaling pathway is presented in [Supplementary Fig. 1](#) (available online). Steps for selecting SNPs are shown in [Fig. 1](#). In terms of gene-

set analysis using MAGMA for genes in the Hedgehog signaling pathway, we identified that *KIF27* harboring 100 SNPs ( $Z=1.88$ ,  $P=0.03$ ; [Fig. 2](#) and [Supplementary Table 3](#), available online) showed the strongest relationship with the gastric cancer susceptibility. After function prediction based on Regulome DB, SNP info Web Server, HaploReg, and LD analysis, only 5 SNPs were kept for further investigation ([Supplementary Table 4](#), available online).

In single-variant genetic association analysis, rs2990912 and rs2065516 were detected to be significantly correlated with gastric cancer susceptibility in the additive model. After FDR correction, only rs2990912 in *KIF27* was related to gastric cancer susceptibility ( $P=2.77\times 10^{-2}$ ; [Table 1](#)).

### Association between SNP rs2990912 of *KIF27* and the risk of gastric cancer


We investigated the significance of the association between SNP rs2990912 with gastric cancer risk using additive, dominant, codominant, and recessive genetic effect models. The frequencies of AA, AG and GG were 83.4%, 16.0%, and 0.6% in gastric cancer cases and 86.6%, 13.1%, and 0.3% in controls, respectively. More individuals carried G allele were observed in cases than in controls compared with the AA genotype in the dominant model (OR, 1.29; 95% CI, 1.07–1.55;  $P=7.39\times 10^{-3}$ , [Table 2](#)), indicating that individuals who carried rs2990912 G allele had a 1.29-fold increased risk of gastric cancer.

Further stratification analyses were performed according to age and gender. The results showed that individuals carried G allele in rs2990912 had a prominently increased gastric cancer risk compared with subjects carried A allele in the subgroups of younger individuals with an age less than 60 years (OR, 1.43; 95% CI, 1.10–1.86;  $P=7.56\times 10^{-3}$ ) and males (OR, 1.27; 95% CI, 1.03–1.56;  $P=7.39\times 10^{-3}$ , [Table 3](#)). We further conducted Kaplan-Meier survival analysis using rs2990912 genotype and clinical data obtained from TCGA to evaluate whether rs2990912 played a role in gastric cancer patients' survival. As shown in [Supplementary Fig. 2](#) (available online), rs2990912 did not change the survival of gastric cancer patients.

### Gene expression in gastric tumor and normal tissues

We evaluated the expression pattern of *KIF27* in gastric tumor and normal tissues using TCGA and GEO datasets, and found that *KIF27* expression in tumors was significantly higher than that in normal





**Fig. 2 Gene-set analysis.** Gene-set analysis using MAGMA for genes in the Hedgehog signaling pathway. The top panel shows the SNP numbers located in genes; and the bottom panel shows the *P*-value of genes. SNPs: single nucleotide polymorphisms; MAGMA: Multi-marker Analysis of GenoMic Annotation.

Chr	SNP	Gene	Allele <sup>a</sup>	MAF (case)	MAF (control)	OR (95% CI) <sup>b</sup>	<i>P</i> <sup>b</sup>	<i>P</i> <sup>c</sup>
9	rs2990912	<i>KIF27</i>	A/G	0.09	0.07	1.29(1.08–1.54)	$5.53 \times 10^{-3}$	$2.77 \times 10^{-2}$
9	rs2065516	<i>KIF27</i>	C/T	0.44	0.47	0.90(0.82–0.99)	$2.84 \times 10^{-2}$	$7.10 \times 10^{-2}$

<sup>a</sup>Reference allele/effect allele; <sup>b</sup>*P*-values for additive model adjusted for sex and age in logistic regression model; <sup>c</sup>*P*-values after false discovery rate correction. Chr: chromosome; MAF: minor allele frequency; OR: odds ratio; CI: confidence interval.

Genotypes	Cases		Controls		OR (95% CI) <sup>a</sup>	<i>P</i> <sup>a</sup>
	<i>N</i>	%	<i>N</i>	%		
AA	1324	83.4	1761	86.6	1.00	
AG	253	16.0	266	13.1	1.27 (1.06–1.54)	$1.16 \times 10^{-2}$
GG	9	0.6	6	0.3	1.95 (0.69–5.51)	$2.10 \times 10^{-1}$
Additive model					1.29 (1.08–1.54)	$5.53 \times 10^{-3}$
Dominant model (AG+GG vs. AA)					1.29 (1.07–1.55)	$7.39 \times 10^{-3}$
Recessive model (GG vs. AG+AA)					1.88 (0.66–5.32)	$2.35 \times 10^{-1}$

<sup>a</sup>ORs and *P*-values were adjusted for age and sex in logistic regression model. OR: odds ratio; CI: confidence interval.


tissues (**Fig. 3**). We observed the expression of *KIF27* in the pathological characteristics of gastric cancer using clinical data obtained from TCGA and found that *KIF27* expression was significant different in tumor stage II and III (**Supplementary Fig. 3**, available online).


Additionally, we assessed the impact of SNPs on the *KIF27* expression in different tumor grades and stages using data obtained from TCGA database and performed the eQTL analysis using the GTEx database. As shown in **Supplementary Fig. 4** (available online), rs2990912 genotype did not affect

**Table 3** Stratified analysis for rs2990912 and gastric cancer risk in additive model

Variables	rs2990912 (cases/controls)						OR (95% CI) <sup>a</sup>	<i>P</i> <sup>a</sup>
	AA		AG		GG			
	<i>N</i>	%	<i>N</i>	%	<i>N</i>	%		
<b>Age (years)</b>								
<60	591/840	83.0/87.1	115/123	16.2/12.8	6/1	0.8/0.1	1.43 (1.10–1.86)	7.56×10 <sup>-3</sup>
≥60	733/921	83.9/86.1	138/143	15.8/13.4	3/5	0.3/0.5	1.17 (0.92–1.49)	1.94×10 <sup>-1</sup>
<b>Sex</b>								
Male	1029/1197	83.8/86.7	191/180	15.6/13.0	8/4	0.6/0.3	1.27 (1.03–1.56)	7.39×10 <sup>-3</sup>
Female	295/564	82.4/86.5	62/86	17.3/13.2	1/2	0.3/0.3	1.34 (0.95–1.89)	9.32×10 <sup>-2</sup>

<sup>a</sup>ORs and *P*-values for additive model adjusted for sex and age in logistic regression model. OR: odds ratio; CI: confidence interval.

**Fig. 3** Expression of *KIF27* in cancer and normal tissues. Differences in *KIF27* expression between normal tissues and gastric tumor tissues based on TCGA database (A and B), GSE66229 (C and D), GSE13911 (E), GSE37023 (F). *N*: number of gastric tumor or normal tissues. *P*-values were calculated using two-sided Mann-Whitney test.



**Fig. 4 Association of *KIF27* expression with patients' survival.** Kaplan-Meier survival curve of gastric cancer patients according to the *KIF27* mRNA status by using Kaplan-Meier Plotter, an online survival analysis tool (<http://kmplot.com/analysis/>) of overall survival (A), progression-free survival (B), and post-progression survival (C).

*KIF27* expression in each grade and stage. Although there was no eQTL of rs2990912 on *KIF27* in gastric tissues (*Supplementary Fig. 5A*, available online), we observed that rs2990912 G allele could increase *KIF27* expression in the tissues of the digestive system including liver tissues (*Supplementary Fig. 5B*, available online;  $P=9.60 \times 10^{-3}$ ) and small intestine tissues (*Supplementary Fig. 5C*, available online;  $P=5.30 \times 10^{-3}$ ). Moreover, the regulation of rs2990912 on the expression of *KIF27* in gastric tissues was consistent with that in other digestive system tissues (*Supplementary Fig. 6*, available online).

We further explored the effect of *H. pylori* infection on the expression level of *KIF27* and found no significant difference (*Supplementary Fig. 7A*, available online). In addition, *KIF27* expression did not change significantly according to the SNP genotype regardless of the presence or absence of *H. pylori* infection (*Supplementary Fig. 7B*, available online).

#### Effect of *KIF27* expression on gastric cancer patient survival

Kaplan-Meier plotter was applied to assess whether *KIF27* expression would affect the survival of gastric cancer patients. Individuals with higher *KIF27* expression had prominently shorter overall survival time (HR, 1.6; 95% CI, 1.24–2.07;  $P=2.80 \times 10^{-4}$ , *Fig. 4A*), worse progression-free survival (HR, 1.47; 95% CI, 1.12–1.92;  $P=4.80 \times 10^{-3}$ , *Fig. 4B*), and poor post-progression survival (HR, 1.7; 95% CI, 1.23–2.35;  $P=1.30 \times 10^{-3}$ , *Fig. 4C*). Further, in the stratified analysis on clinicopathological features, we observed that the effects of *KIF27* expression on survival were significant in the subgroups of male, stage T2, stage T4, stage N1, stage M0, Lauren

classification, and differentiation (*Supplementary Table 5*, available online).

#### Association between *KIF27* expression and immune infiltrates

We next investigated the relationship between *KIF27* expression and immune cell infiltration degree in gastric cancer via the TIMER2.0 web server. The results showed that *KIF27* expression was significantly associated with the infiltration of several immune cells, including CD8<sup>+</sup> T cells (Rho=0.222,  $P=1.34 \times 10^{-5}$ ), CD4<sup>+</sup> T cells (Rho=0.398,  $P=8 \times 10^{-16}$ ), macrophages (Rho=0.144,  $P=5.13 \times 10^{-3}$ ), Tregs (Rho=0.295,  $P=4.93 \times 10^{-9}$ ), and neutrophils (Rho=0.196,  $P=1.19 \times 10^{-4}$ , *Supplementary Fig. 8*, available online).

The relationship between *KIF27* expression and the expression level of immune markers was further explored. As shown in *Supplementary Fig. 9* (available online), *KIF27* expression was related to the expression level of several immune markers, including those specific to monocytes, TAM, M1 macrophage, and M2 macrophage. In addition, the result of correlation analysis showed that *KIF27* expression was related to PD-L1 expression (Rho=0.24,  $P=3.2 \times 10^{-9}$ , *Supplementary Fig. 10*, available online).

#### Discussion

The Hedgehog signaling pathway contributes to tumorigenesis. Abnormal activation of the Hedgehog signaling pathway causes the destruction of gastric cell differentiation, loss of gastric acid secretion, and tumor transformation<sup>[24]</sup>. Dysregulation of the Hedgehog signaling leads to the invasion and



metastasis of gastric cancer and the obstruction of it reduces the proliferation of tumor cells<sup>[25]</sup>. The signaling molecules, such as *SHH*, *IHH*, *PTCH1*, and *GLII*, are used as biomarkers in clinical oncology. The small molecule compounds targeting SMO and humanized anti-SHH antibodies are considered to be effective anti-cancer drugs for gastric cancer<sup>[12]</sup>. Previous studies have shown that genetic variants in the Hedgehog signaling pathway genes were related to the occurrence and development of diseases. The polymorphism of the *GLII* gene SNP rs2228226 was closely related to the risk of chronic lymphocytic leukemia in the Chinese population<sup>[26]</sup>. *GLI3* genetic variants were associated with overall bladder cancer risk, and two *GLI2* SNPs and one *SHH* SNP were significantly associated with the overall survival of patients with muscle-invasive and metastatic bladder cancer<sup>[19]</sup>. The GG genotype of the *HHIP* genetic variant rs1489759 had a protective effect on chronic obstructive pulmonary disease and lung cancer<sup>[20]</sup>. Most of these studies focused on a few key genes in the Hedgehog signaling pathway, such as *GLI*, *HH*, and *HHIP*. Yet, the association between the genetic variation with gastric cancer susceptibility has rarely been reported. Although genetic variants of *GLII* were associated with inflammatory bowel diseases susceptibility<sup>[27]</sup>, we did not find a correlation between genetic variants of these reported genes and the risk of gastric cancer. Given that the Hedgehog signaling pathway plays an important role in gastrointestinal tract development, homeostasis, and malignancy, we continued to evaluate the association of the genetic variants of 31 Hedgehog signaling pathway-related genes with gastric cancer susceptibility. We found that the G allele in *KIF27* was correlated with a higher gastric cancer risk, especially in the younger and male subjects.

*KIF27*, kinesin family member 27, is located at human chromosome 9q22.1. *KIF27* and *KIF7* are homologs of *Drosophila* *Cos2* in human. *KIF27* is most homologous to *Cos2*, sharing the common domain structure with *Drosophila* *Cos2*, while *KIF7* shares only part of domain structure with *Drosophila* *Cos2*<sup>[28–30]</sup>. In *Drosophila*, *Fu* (*STK36*) and *Cos2* regulate the cell surface accumulation and signal transduction activity of SMO by regulating SMO phosphorylation. *Cos2* interacts with SMO to prevent HH-induced SMO phosphorylation, while *Fu* promotes SMO phosphorylation by antagonizing *Cos2*<sup>[31]</sup>. SMO activity plays a very important role in

the downstream cascade activation by blocking GLIR and promoting GLIA<sup>[32]</sup>. Despite the conflicting reports about the role of *KIF7* and *KIF27* in Hedgehog signaling in vertebrates<sup>[33–34]</sup>, *KIF27* retains its partnership with *Fu* and participates in the recruitment and/or activation of proteins along the center of the cilia<sup>[35]</sup>. Evidence showed that low-level expression of *KIF7* was associated with the poor prognosis of epithelial ovarian cancer and *KIF7* inhibited prostate cancer development as a negative regulator<sup>[36]</sup>. It is reported that *KIF27* was involved in leukemogenesis and was a component of a minimal deleted region (MDR) in acute myeloid leukemia<sup>[37]</sup>. There has been no report about the association of *KIF27* with gastric cancer. We observed *KIF27* was significantly increased in gastric tumors compared with normal tissues. *KIF27* may be a participant in the occurrence and development of gastric cancer.

Evidence has shown that *SHH* is a gastric morphogen and can initiate gastritis in response to *H. pylori* infection<sup>[38]</sup>. Studies have reported that the Hedgehog signaling pathway regulates the expression of PD-L1 and may affect the anti-tumor activity of lymphocytes<sup>[39]</sup>. Furthermore, data have suggested HH/GLI inhibitor can drastically reduce PD-L1 expression and inhibit the proliferation of tumor cells in gastric cancer<sup>[40]</sup>. Given the important role of this signal in cancers, we assessed the potential impact of *KIF27* on the survival of patients with gastric cancer and found that elevated *KIF27* was related to poor gastric outcomes. Furthermore, elevated *KIF27* expression in gastric cancer was related to a higher N stage HR in OS (*Supplementary Table 5*). The results showed that *KIF27* expression was associated with immune infiltration degree in gastric cancer. We further found that *KIF27* expression was related to marker expression of different immune cell subsets. These results suggested *KIF27* may interact within the tumor microenvironment. Although our results showed that *H. pylori* infection did not affect the expression pattern of *KIF27*, we found *KIF27* was positively correlated with *PD-L1*. Thus, further functional studies are needed to determine the biological mechanisms of *KIF27* and genetic variants involved in the occurrence and development of gastric cancer.

In summary, we identified a genetic variant rs2990912 in *KIF27*, which may influence gastric cancer susceptibility in the Chinese Han population. *KIF27* and the genetic variants in it are potential

biomarkers for the assessment of gastric cancer risk and may hold important value for the diagnosis, prognosis, and therapy of gastric cancer.

### Acknowledgments

This study was funded by the National Key R&D Program of China (Grants No. 2018YFC1313100 and No. 2018YFC1313102) and the National Natural Science Foundation of China (Grants No. 81773538 and No. 81773539), Collaborative Innovation Center for Cancer Personalized Medicine, and the Priority Academic Program Development of Jiangsu Higher Education Institutions (Public Health and Preventive Medicine). We appreciate comments from editor and anonymous reviewers.

### References

- [1] Bray F, Ferlay J, Soerjomataram I, et al. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries[J]. *CA Cancer J Clin*, 2018, 68: 394–424.
- [2] Zheng R, Zeng H, Zhang S, et al. Estimates of cancer incidence and mortality in China, 2013[J]. *Chin J Cancer*, 2017, 36: 66.
- [3] Hunt RH, Camilleri M, Crowe SE, et al. The stomach in health and disease[J]. *Gut*, 2015, 64: 1650–1668.
- [4] Mayer RJ, Venook AP, Schilsky RL. Progress against GI cancer during the American Society of Clinical Oncology's first 50 years[J]. *J Clin Oncol*, 2014, 32: 1521–1530.
- [5] Labreche K, Daniau M, Sud A, et al. A genome-wide association study identifies susceptibility loci for primary central nervous system lymphoma at 6p25.3 and 3p22.1: a LOC network study[J]. *Neuro Oncol*, 2019, 21(8): 1039–1048.
- [6] Hu N, Wang Z, Song X, et al. Genome-wide association study of gastric adenocarcinoma in Asia: a comparison of associations between cardia and non-cardia tumours[J]. *Gut*, 2016, 65: 1611–1618.
- [7] Yang Y, Basu S, Zhang L. A Bayesian hierarchical variable selection prior for pathway-based GWAS using summary statistics[J]. *Stat Med*, 2020, 39: 724–739.
- [8] Gu D, Li S, Ben S, et al. Circadian clock pathway genes associated with colorectal cancer risk and prognosis[J]. *Arch Toxicol*, 2018, 92: 2681–2689.
- [9] Tang D, Liu H, Zhao Y, et al. Genetic variants of BIRC3 and NRG1 in the NLRP3 inflammasome pathway are associated with non-small cell lung cancer survival[J]. *Am J Cancer Res*, 2020, 10: 2582–2595.
- [10] Lum L, Beachy PA. The Hedgehog response network: sensors, switches, and routers[J]. *Science*, 2004, 304: 1755–1759.
- [11] Huang S, Zhang Z, Zhang C, et al. Activation of Smurf E3 ligase promoted by smoothed regulates hedgehog signaling through targeting patched turnover[J]. *PLoS Biol*, 2013, 11: e1001721.
- [12] Katoh Y, Katoh M. Hedgehog signaling pathway and gastrointestinal stem cell signaling network (review)[J]. *Int J Mol Med*, 2006, 18: 1019–1023.
- [13] Katoh M. Genomic testing, tumor microenvironment and targeted therapy of Hedgehog-related human cancers[J]. *Clin Sci (Lond)*, 2019, 133: 953–970.
- [14] Saqui-Salces M, Coves-Datson E, Veniaminova NA, et al. Inflammation and Gli2 suppress gastrin gene expression in a murine model of antral hyperplasia[J]. *PLoS One*, 2012, 7: e48039.
- [15] Song Y, Tu J, Cheng Y, et al. HHIP Overexpression Suppresses Human Gastric Cancer Progression and Metastasis by Reducing Its CpG Island Methylation[J]. *Front Oncol*, 2020, 10: 1667.
- [16] Peng Y, Zhang X, Lin H, et al. Dual activation of Hedgehog and Wnt/beta-catenin signaling pathway caused by downregulation of SUFU targeted by miRNA-150 in human gastric cancer[J]. *Aging (Albany NY)*, 2021, 13: 10749–10769.
- [17] Koh V, Chakrabarti J, Torvund M, et al. Hedgehog transcriptional effector GLI mediates mTOR-Induced PD-L1 expression in gastric cancer organoids[J]. *Cancer Lett*, 2021, 518: 59–71.
- [18] Cao W, Li Y, Sun H, et al. Apatinib Suppresses Gastric Cancer Stem Cells Properties by Inhibiting the Sonic Hedgehog Pathway[J]. *Front Cell Dev Biol*, 2021, 9: 679806.
- [19] Chen M, Hildebrandt MA, Clague J, et al. Genetic variations in the sonic hedgehog pathway affect clinical outcomes in non-muscle-invasive bladder cancer[J]. *Cancer Prev Res (Phila)*, 2010, 3: 1235–1245.
- [20] Young RP, Whittington CF, Hopkins RJ, et al. Chromosome 4q31 locus in COPD is also associated with lung cancer[J]. *Eur Respir J*, 2010, 36: 1375–1382.
- [21] Howie BN, Donnelly P, Marchini J. A flexible and accurate genotype imputation method for the next generation of genome-wide association studies[J]. *PLoS Genet*, 2009, 5: e1000529.
- [22] Abnet CC, Freedman ND, Hu N, et al. A shared susceptibility locus in PLCE1 at 10q23 for gastric adenocarcinoma and esophageal squamous cell carcinoma[J]. *Nat Genet*, 2010, 42: 764–767.
- [23] de Leeuw CA, Mooij JM, Heskes T, et al. MAGMA: generalized gene-set analysis of GWAS data[J]. *PLoS Comput Biol*, 2015, 11: e1004219.
- [24] Konstantinou D, Bertaux-Skeirik N, Zavros Y. Hedgehog signaling in the stomach[J]. *Curr Opin Pharmacol*, 2016, 31: 76–82.

- [25] Skoda AM, Simovic D, Karin V, et al. The role of the Hedgehog signaling pathway in cancer: A comprehensive review[J]. *Bosn J Basic Med Sci*, 2018, 18: 8–20.
- [26] Li YY, Tian T, Zhang R, et al. Association between polymorphism of *GLI1* gene SNP rs2228226 and chronic lymphocytic leukemia in Chinese population[J]. *Med Oncol*, 2014, 31: 294.
- [27] Lees CW, Zacharias WJ, Tremelling M, et al. Analysis of germline *GLI1* variation implicates hedgehog signalling in the regulation of intestinal inflammatory pathways[J]. *PLoS Med*, 2008, 5: e239.
- [28] Endoh-Yamagami S, Evangelista M, Wilson D, et al. The mammalian *Cos2* homolog *Kif7* plays an essential role in modulating Hh signal transduction during development[J]. *Curr Biol*, 2009, 19: 1320–1326.
- [29] Katoh Y, Katoh M. Characterization of *KIF7* gene in silico[J]. *Int J Oncol*, 2004, 25: 1881–1886.
- [30] Katoh Y, Katoh M. *KIF27* is one of orthologs for *Drosophila Costal-2*[J]. *Int J Oncol*, 2004, 25: 1875–1880.
- [31] Liu Y, Cao X, Jiang J, et al. Fused-Costal2 protein complex regulates Hedgehog-induced Smo phosphorylation and cell-surface accumulation[J]. *Genes Dev*, 2007, 21: 1949–1963.
- [32] Szczepny A, Rogers S, Jayasekara WSN, et al. The role of canonical and non-canonical Hedgehog signaling in tumor progression in a mouse model of small cell lung cancer[J]. *Oncogene*, 2017, 36: 5544–5550.
- [33] Tay SY, Ingham PW, Roy S. A homologue of the *Drosophila* kinesin-like protein *Costal2* regulates Hedgehog signal transduction in the vertebrate embryo[J]. *Development*, 2005, 132: 625–634.
- [34] Varjosalo M, Li SP, Taipale J. Divergence of hedgehog signal transduction mechanism between *Drosophila* and mammals[J]. *Dev Cell*, 2006, 10: 177–186.
- [35] Wilson CW, Nguyen CT, Chen MH, et al. Fused has evolved divergent roles in vertebrate Hedgehog signalling and motile ciliogenesis[J]. *Nature*, 2009, 459: 98–102.
- [36] Yao Y, Liu L, He W, et al. Low expression of *KIF7* indicates poor prognosis in epithelial ovarian cancer[J]. *Cancer Biomark*, 2019, 26: 481–489.
- [37] Naarmann-de Vries IS, Sackmann Y, Klein F, et al. Characterization of acute myeloid leukemia with *del(9q)* - Impact of the genes in the minimally deleted region[J]. *Leuk Res*, 2019, 76: 15–23.
- [38] Schumacher MA, Feng R, Aihara E, et al. *Helicobacter pylori*-induced Sonic Hedgehog expression is regulated by NFkappaB pathway activation: the use of a novel in vitro model to study epithelial response to infection[J]. *Helicobacter*, 2015, 20: 19–28.
- [39] Onishi H, Fujimura A, Oyama Y, et al. Hedgehog signaling regulates PDL-1 expression in cancer cells to induce anti-tumor activity by activated lymphocytes[J]. *Cell Immunol*, 2016, 310: 199–204.
- [40] Chakrabarti J, Holokai L, Syu L, et al. Hedgehog signaling induces PD-L1 expression and tumor cell proliferation in gastric cancer[J]. *Oncotarget*, 2018, 9: 37439–37457.

Submit to the *Journal* by ScholarOne Manuscripts at  
<http://mc03.manuscriptcentral.com/jbrint>

